

## CO-RANK AND BETTI NUMBER OF A GROUP

IRINA GELBUKH, Mexico City

(Received December 24, 2014)

*Abstract.* We study the maximal ranks of a free and a free abelian quotients of a finitely generated group, called co-rank (inner rank, cut number) and the Betti number, respectively. We show that any combination of these values within obvious constraints is realized for some finitely presented group, which is important for manifold and foliation topology.

*Keywords:* co-rank, inner rank, fundamental group

*MSC 2010:* 20E05, 20F34, 14F35

We study the relation between the co-rank  $\text{corank}(G)$  of a finitely generated group  $G$  and its Betti number  $b(G)$ . These values bound the isotropy index  $i(G)$  of  $G$ :  $\text{corank}(G) \leq i(G) \leq b(G)$  [2, 3, 11]. These notions have important applications in theory of manifolds as the first non-commutative Betti number  $b'_1(M) = \text{corank}(\pi_1(M))$ , the first Betti number  $b_1(M) = b(\pi_1(M))$ , and  $h(M) = i(\pi_1(M))$ , where  $\pi_1(M)$  is the fundamental group of the manifold  $M$ . For any  $n \geq 4$ , a group is the fundamental group of a smooth closed connected  $n$ -manifold iff it is finitely presented. In theory of 2- and 3-manifolds, co-rank of the fundamental group coincides with the cut-number, a generalization of the genus for closed surfaces [7, 13]. In the theory of foliations of Morse forms,  $b'_1(M)$  and  $h(M)$  define the topology of the foliation [3, 4], the form's cohomology class [5], and the types of its singularities [6].

For a finitely generated abelian group  $G = \mathbb{Z}^n \oplus T$ , where  $T$  is finite, its *torsion-free rank*, *Prüfer rank*, or *(first) Betti number*, is  $b(G) = \text{rk}(G/T) = n$ . The latter term extends to finitely generated groups by  $b(G) = b(G^{ab}) = \text{rk}(G^{ab}/T(G^{ab}))$ , where  $G^{ab} = G/[G, G]$  is the abelianization and  $T(\cdot)$ , the torsion subgroup. In other words:

**Definition 1.** The Betti number  $b(G)$  of a finitely generated group  $G$  is the maximum rank of a free abelian quotient group of  $G$ , i.e., the maximum rank of a free abelian group  $A$  such that there exists an epimorphism  $\varphi : G \twoheadrightarrow A$ .

The term came from geometric group theory, where  $G^{ab}$  is called the first homology group  $H_1(G)$ . A non-commutative analog of Betti number can be defined as follows:

**Definition 2** ([7, 8]). *The co-rank  $\text{corank}(G)$  [8], inner rank  $IN(G)$  [7] or  $Ir(G)$  [9], or first non-commutative Betti number  $b'_1(G)$  [1] of a finitely generated group  $G$  is the maximum rank of a free quotient group of  $G$ , i.e., the maximum rank of a free group  $F$  such that there exists an epimorphism  $\varphi : G \twoheadrightarrow F$ .*

The notion of co-rank is also in a way dual to that of rank, which is the minimum rank of a free group allowing an epimorphism onto  $G$ . In contrast to rank, co-rank is algorithmically computable for finitely presented groups [10, 12].

For example,  $\text{corank}(\mathbb{Z}^n) = 1$ , while  $b(\mathbb{Z}^n) = n$ . For a finite group  $G$ ,  $\text{corank}(G) = b(G) = 0$ ; the same holds for  $G = \mathbb{Z}_2 * \mathbb{Z}_2 * \mathbb{Z}_2$ , even though it is infinite and contains  $F_2$  and thus free subgroups of all ranks up to countable. Obviously, for any finitely generated group,  $\text{corank}(G) \leq b(G) \leq \text{rk } G$  and  $b(G) \geq 1$  implies  $\text{corank}(G) \geq 1$ . In this paper we show that these are the only constraints between these values:

**Theorem 3.** *Let  $0 \leq c, b, r \in \mathbb{Z}$ . Then there exists a finitely generated group  $G$  with  $\text{corank}(G) = c$ ,  $b(G) = b$ , and  $\text{rk } G = r$  iff*

$$c = b = 0 \quad \text{or} \quad 1 \leq c \leq b \leq r;$$

*the group can be chosen to be finitely presented and, if  $b = r$ , torsion-free.*

**Lemma 4.** *Let  $G_1, G_2$  be finitely generated groups. Then for the Betti number of the free product and of the direct product,*

$$b(G_1 * G_2) = b(G_1 \times G_2) = b(G_1) + b(G_2).$$

*Proof.* Obviously,  $(G_1 * G_2)^{ab} = (G_1 \times G_2)^{ab}$ . Denote  $G = G_1 \times G_2$ . Since epimorphisms  $G_i \twoheadrightarrow \mathbb{Z}^{b(G_i)}$  onto free abelian groups can be extended to an epimorphism of  $G_1 \times G_2 \twoheadrightarrow \mathbb{Z}^{b(G_2)} \times \mathbb{Z}^{b(G_2)} = \mathbb{Z}^{b(G_2)+b(G_2)}$ , we have  $b(G) \geq b(G_1) + b(G_2)$ .

Let us now show that  $b(G) \leq b(G_1) + b(G_2)$ . Consider the natural homomorphisms  $\psi_1 : G_1 \rightarrow G_1 \times 1 \subseteq G$ ,  $\psi_2 : G_2 \rightarrow 1 \times G_2 \subseteq G$ . Then  $\psi_i$  and an epimorphism onto a free abelian group

$$G_i \xrightarrow{\psi_i} G = G_1 \times G_2 \twoheadrightarrow A = \mathbb{Z}^{b(G)}$$

induces a homomorphism  $\varphi_i : G_i \rightarrow A$ . Since  $A_i = \varphi_i(G_i) \subseteq A$  are free abelian groups,  $\text{rk } A_i \leq b(G_i)$ . Since  $G = \langle \psi_1(G_1), \psi_2(G_2) \rangle$ , we have  $A = \langle A_1, A_2 \rangle$ ; in particular,  $b(G) = \text{rk } A \leq \text{rk } A_1 + \text{rk } A_2$ .  $\square$

*Proof of Theorem 3.* For  $1 \leq c \leq b \leq r$ , consider  $G = \mathbb{Z}^{b_1} * \dots * \mathbb{Z}^{b_c} * \mathbb{Z}_2^{r-b}$  such that  $\sum_{i=1}^c b_i = b$ . By [9, Proposition 6.4],  $\text{corank}(G_1 * G_2) = \text{corank}(G_1) + \text{corank}(G_2)$ , so  $\text{corank}(G) = \sum_{i=1}^c \text{corank}(\mathbb{Z}^{b_i}) = c$ . By Lemma 4,  $b(G) = \sum_{i=1}^c b(\mathbb{Z}^{b_i}) = b$ , and by Grushko-Neumann theorem,  $\text{rk } G = r$ .  $\square$

# REFERENCES

- [1] P. Arnoux, G. Levitt: Sur l'unique ergodicité des 1-formes fermées singulières, *Invent. Math.* 84 (1986), 141–156. Zbl 0577.58021, MR0830042
- [2] A. Dimca, S. Papadima, A. Suciu: Quasi-Kähler groups, 3-manifold groups, and formality, *Mathematische Zeitschrift* 268 (2011), no. 1–2, 169–186. Zbl 1228.14018, MR2805428
- [3] I. Gelbukh: Number of minimal components and homologically independent compact leaves for a Morse form foliation, *Stud. Sci. Math. Hung.* 46 (2009), no. 4, 547–557. Zbl 1274.57005, MR2654204
- [4] I. Gelbukh: On the structure of a Morse form foliation, *Czech. Math. J.* 59 (2009), no. 1, 207–220. Zbl 1224.57010, MR2486626
- [5] I. Gelbukh: Close cohomologous Morse forms with compact leaves, *Czech. Math. J.* 63 (2013), no. 2, 515–528. Zbl 1289.57009, MR3073975
- [6] I. Gelbukh: The number of split points of a Morse form and the structure of its foliation, *Math. Slovaca* 63 (2013), no. 2, 331–348. Zbl 06163523, MR3037071
- [7] W. Jaco: Geometric realizations for free quotients, *J. Austral. Math. Soc.* 14 (1972), 411–418. Zbl 0259.57004, MR0316571
- [8] C. J. Leininger, A. W. Reid: The co-rank conjecture for 3-manifold groups, *Algebraic and Geometric Topology* 2 (2002), 37–50. Zbl 0983.57001, MR1885215
- [9] R. C. Lyndon, P. E. Schupp: *Combinatorial group theory*, Mathematics, Springer, Berlin, 2001. Zbl 0997.20037, MR1812024
- [10] G. S. Makanin: Equations in a free group, *Math. USSR Izvestiya* 21 (1983), no. 3, 483–546. Zbl 0527.20018, MR0682490
- [11] I. A. Mel'nikova: Maximal isotropic subspaces of skew-symmetric bilinear mapping, *Mosc. Univ. Math. Bull.* 54 (1999), no. 4, 1–3. Zbl 0957.57018, MR1716286
- [12] A. A. Razborov: Systems of equations in a free group, *Math. USSR Izvestiya* 25 (1985), 115–162. Zbl 0579.20019, MR0755958
- [13] A. Sikora: Cut numbers of 3-manifolds, *Trans. Amer. Math. Soc.* 357 (2005), no. 5, 2007–2020. Zbl 1064.57018, MR2115088

*Authors' addresses:* Irina Gelbukh, Centro de Investigación en Computación (CIC), Instituto Politécnico Nacional (IPN), Av. Juan de Dios Bátiz, 07738, DF, Mexico City, Mexico, e-mail: [gelbukh@member.ams.org](mailto:gelbukh@member.ams.org).